Appl No. 10/051,401 Amdt. Dated October 21, 2004 Reply to Office Action of July 21, 2004 Customer No. 27752

This listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

- 1. (Currently Amended) An anhydrous treatment composition comprising:
 - a) a reactive agent, wherein said reactive agent additionally comprises a cosmetically active functional group, and
 - b) a solvent in which the reactive agent is soluble and which is water-miscible.
- 2. An anhydrous treatment composition according to Claim 1 which comprises from about 0.01% to about 10%, by weight of the composition, of the reactive agent, and from about 25% to about 95%, by weight of the composition, of the water-miscible solvent.
- 3. An anhydrous treatment composition according to Claim 2, wherein the reactive agent is selected from the group consisting of electrophilic groups, nucleophilic groups or protected thiol groups.
- 4. (Canceled)
- 5. (Currently Amended) An anhydrous treatment composition according to Claim [[4]]1, wherein the reactive agent comprises an electrophilic reactive group selected from the group consisting of halotriazines, haloquinoxalines, halopyrimidines, vinylsulfones, β-haloethylsulfones, β-sulfatoethylsulfones, acrylates, methacrylates, acrylamides, methacrylamides, maleimides, epoxides, acylhalides, esters, carbamates, dithiocarboxylic acid esters, alkoxysilanes, thiosulfates, anhydrides, urea derivatives, isothiocyanates, isocyanates, lactones, thiosulfates, isothiuroniums, azalactones and mixtures thereof.
- 6. (Currently Amended) An anhydrous treatment composition according to Claim [[4]]1, wherein the reactive agent comprises a nucleophilic reactive group selected from the group consisting of thiols, thiolates, thiols or thiolates containing quaternary salts, thioalkyl esters, thioalkylamides, thiol or thiolate derivatives of cysteamine, and mixtures thereof.
- 7. (Currently Amended) An anhydrous treatment composition according to Claim [[4]]], wherein the reactive agent comprises a protected thiol reactive group having the formula

Appl No. 10/051,401 Amdr. Dated October 21, 2004 Reply to Office Action of July 21, 2004 Customer No. 27752

$$R-(S-Pr)_m$$

where R is a mono or multivalent cosmetically active functional group, S is sulfur, Pr is a protecting group and m is an integer between 1 and 100.

- 8. An anhydrous treatment composition according to Claim 7, wherein the protecting group is selected from the group consisting of heterocyclic protecting groups, sp² aliphatic trigonal carbon protecting groups, sp³ carbon electrophilic protecting groups, phosphorus protecting groups, metal based protecting groups, non-metal and metalloid based protecting groups other than phosphorus, energy-sensitive protecting groups and mixtures thereof.
- An anhydrous treatment composition according to Claim 1, wherein the water-miscible solvent has a Vaughan Solubility Parameter of from about 8.0 (cal/cm³)^{0.5} to about 17.0 (cal/cm³)^{0.5}.
- An anhydrous treatment composition according to Claim 1, wherein the water-miscible 10. solvent is selected from the group consisting of ethyl formate, dimethyl isosorbide, acetylacetone, 2-butanone, acetone, methyl acetate, ethyl acetate, propyl acetate, ethoxyethanol, dipropylene glycol monomethyl ether, butyl lactate, t-butyl alcohol, phenyl acetate, 2-propoxyethanol, 2-isopropoxyethanol, methoxypropanol, isopropyl lactate, hexyl alcohol, butoxyethanol, tripropylcne glycol (PPG-3), triacetin, methoxyethanol, isopropyl alcohol, PEG-8, methyl lactate, PEG-6, PEG-5, PEG-4, N-methylpyrrolidone, propyl alcohol, dipropylene glycol (PPG-2), acetonitrile, phenoxyethanol, triethylene glycol, hexylene glycol, cthyl alcohol, γ-butyrolactone, butylene glycol, propylene carbonate, dimethyl sulfoxide, dicthylene glycol, ethoxydiglycol, propylene glycol, pyrrolidone, pyrrolidone-2, methyl alcohol, ethylene carbonate, ethylene glycol, acetamide, glycerin, butyl carbitol, 1,3-dioxolane, dimethoxymethane, 1,2-hexanediol, dipropylene glycol butyl ether, dipropylene glycol t-butyl ether, propionaldehyde, diethoxymethane, glycerol formal, y-valerolactone, α-methyl-y-butyrolactone, and butyl acetate, cthanol, isopropyl alcohol, and mixtures thereof.
- 1011. (Currently Amended) An anhydrous treatment according to Claim 910, wherein the water-miscible solvent is selected from the group consisting of N-methylpyrrolidinone, propylene carbonate, γ-butyrolactone, γ-valerolactone, propylene glycol, dipropylene glycol,

Appl No. 10/051,401 Amdt. Dated October 21, 2004 Reply to Office Action of July 21, 2004 Customer No. 27752

ethoxydiglycol, ethoxyethanol, dimethoxymethane, dimethyl isosorbide, butyl lactate, and mixtures thercof.